

Volunteer Lake Assessment Program Individual Lake Reports BEECH POND, LOWER, TUFTONBORO, NH

MORPHOMETRIC DATA TROPHIC CLASSIFICATION KNOWN EXOTIC SPECIES

Watershed Area (Ac.):	1,600	Max. Depth (m):	15.2	Flushing Rate (yr1)	0.8	Year	Trophic class	
Surface Area (Ac.):	155	Mean Depth (m):	6.8	P Retention Coef:	0.63	1980	OLIGOTROPHIC	
Shore Length (m):	4,700	Volume (m³):	4,250,500	Elevation (ft):	968	2001	OLIGOTROPHIC	

The Waterbody Report Card tables are generated from the 2012 305(b) report on the status of N.H. waters, and are based on data collected from 2001-2011.

Designated Use	Parameter	Category	Comments				
Aquatic Life	Phosphorus (Total)	Good	>/=5 samples and median is < threshold but > 1/2 threshold value.				
	рН	Slightly Bad	>10% of samples exceed criteria by a small margin (minimum of 2 exceedances).				
	D.O. (mg/L)	Encouraging	< 10 samples and no exceedance of criteria. More data needed.				
	D.O. (% sat)	Encouraging	< 10 samples and no exceedance of criteria. More data needed.				
	Chlorophyll-a	Good	>/=5 samples and median is < threshold but > 1/2 threshold value.				
Primary Contact Recreation	E. coli	Very Good	All bacteria samples <75% of geometric mean criteria, but not enough to calculate geometric mean. Or, all bacte				
			samples are < single sample criteria and calculated Geometric means are less than geometric mean criteria.				
	Chlorophyll-a	Good	At least 10 samples with 1 sample but < 10% of samples exceeding criteria.				

BEACH PRIMARY CONTACT ASSESSMENT STATUS

LO	OWER BEECH POND - WILLIAM LAWRENCE	E. coli	Good	Geometric means < criteria; however at least 1 exceedance of the single sample criteria occurred.			
C	AMP BEACH						

WATERSHED LAND USE SUMMARY

Fry, J., Xian, G., Jin, S., Dewitz, J., Homer, C., Yang, L., Barnes, C., Herold, N., and Wickham, J., 2011. Completion of the 2006 National Land Cover Database

for the Conterminous United States, PERS, Vol. 77(9):858-864. For larger image contact NHDES.



Land Cover Category	% Cover	Land Cover Category	% Cover	Land Cover Category	% Cover
Open Water 17.6		Barren Land	0	Grassland/Herbaceous	0
Developed-Open Space	2.37	Deciduous Forest	7.5	Pasture Hay	0.42
Developed-Low Intensity 0.11		Evergreen Forest	15.44	Cultivated Crops	0.31
Developed-Medium Intensity	0	Mixed Forest	53.72	Woody Wetlands	1.57
Developed-High Intensity 0		Shrub-Scrub	0.44	Emergent Wetlands	0.59



VOLUNTEER LAKE ASSESSMENT PROGRAM INDIVIDUAL LAKE REPORTS LOWER BEECH POND, TUFTONBORO, NH 2013 DATA SUMMARY

Observations and Recommendations (Refer to Table 1 and Historical Deep Spot Data Graphics)

- CHLOROPHYLL-A: Chlorophyll levels were low on each sampling event and much less than the state median. Historical trend analysis indicates relatively stable chlorophyll with high variability between years.
- CONDUCTIVITY/CHLORIDE: Conductivity and chloride values were low throughout the summer. Historical trend analysis indicates relatively stable epilimnetic conductivity with moderate variability between years.
- E. cou: E. coli levels were very low and much less than the state standards for public beaches and surface waters.
- TOTAL PHOSPHORUS: Epilimnetic (upper water layer) and metalimnetic (middle water layer) phosphorus levels were low and much less than the state median. Historical trend analysis indicates relatively stable epilimnetic phosphorus with high variability between years. Hypolimnetic phosphorus was elevated in August potentially due to the release of phosphorus from bottom sediments when dissolved oxygen levels are less than 1.0 mg/L. Inlet and Outlet phosphorus levels were low in August.
- TRANSPARENCY: Transparency was good and better than the state median. Historical trend analysis indicates relatively stable transparency with moderate variability between years.
- Turbidity: Deep spot and tributary turbidity was low on each sampling event.
- PH: Metalimnetic and hypolimnetic pH levels were less than desirable range of 6.5 8.0 units. Historical trend analysis indicates relatively stable epilimnetic pH with moderate variability between years.
- **RECOMMENDED ACTIONS:** Overall water quality was good and trends were stable. The increased frequency and intensity of storm events highlights the importance of managing stormwater runoff from lake and watershed properties, dirt/gravel road, and steep slopes. DES' "Homeowner's Guide to Stormwater Management" is a good resource for lake residents. Keep up the great work!

	Table 1. 2013 Average Water Quality Data for LOWER BEECH POND								
	Alk.	Chlor-a	Chloride	Cond.	E. Coli	Total P	Trans.	Turb.	рН
Station	mg/l	ug/l	mg/l	uS/cm	#/100ml	ug/l	m	ntu	
							NVS		
Epilimnion	3.30	2.08	5	31.9		5	4.33	0.43	6.53
Metalimnion				31.9		7		0.52	6.47
Hypolimnion				28.5		13		0.60	5.72
First Beach					4				
Inlet				29.0	10	3		0.35	6.56
Outlet				31.6	10	3		0.34	6.72
Second Beach				27.2	3			0.28	6.76

NH Median Values: Median values for specific parameters generated from historic lake monitoring

data.

Alkalinity: 4.9 mg/L

Chlorophyll-a: 4.58 mg/m³ Conductivity: 40.0 uS/cm

Chloride: 4 mg/L

Total Phosphorus: 12 ug/L **Transparency:** 3.2 m

pH: 6.6

NH Water Quality Standards: Numeric criteria for specific parameters. Results exceeding criteria are considered a water quality violation.

Chloride: < 230 mg/L (chronic)

E. coli: > 88 cts/100 mL – public beach E. coli: > 406 cts/100 mL – surface waters Turbidity: > 10 NTU above natural level pH: 6.5-8.0 (unless naturally occurring)

HISTORICAL WATER QUALITY TREND ANALYSIS

Parameter	Trend	Explanation	Parameter	Trend	Explanation
pH	Stable	Trend not significant; data moderately variable.	Chlorophyll-a	Stable	Trend not significant; data highly variable.
Conductivity	Stable	Trend not significant; data show low variability.	Transparency	Stable	Trend not significant; data moderately variable.
	•		Phosphorus (epilimnion)	Stable	Trend not significant; data highly variable.



